

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES


Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problems Mailbox.**



(11) (21) (C) **2,001,667**
(22) 1989/10/27
(43) 1990/04/28
(45) 2000/01/04

(72) Matsuda, Hajime, JP

(72) Ito, Kenzo, JP

(72) Taki, Akio, JP

(72) Uejima, Osamu, JP

(73) Shiseido Company Ltd., JP

(51) Int.Cl.⁶ A61K 7/02, A61K 7/48

(30) 1988/10/28 (63-272653) JP

(54) **COMPOSE COSMETIQUE CONTENANT UN PRODUIT
D'INCLUSION AVEC DE LA CYCLODEXTRINE
HYDROXYALKYLEE**

(54) **COSMETIC COMPOSITION CONTAINING INCLUSION
PRODUCT WITH HYDROXYALKYLATED CYCLODEXTRIN**

(57) A cosmetic composition comprising an inclusion product having a slightly water-soluble component with a hydroxyalkylated cyclodextrin formulated therein.



ABSTRACT OF THE DISCLOSURE

A cosmetic composition comprising an inclusion product having a slightly water-soluble component with a hydroxyalkylated cyclodextrin formulated therein.

- 1 -

COSMETIC COMPOSITION CONTAINING INCLUSION PRODUCT
WITH HYDROXYALKYLATED CYCLODEXTRIN

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates to a cosmetic composition including a liquid and powdery cosmetic compositions comprising an inclusion product having a slightly water-soluble component with a hydroxy-alkylated cyclodextrin (hereinafter abbreviated as "HACD") formulated therein. More specifically, the present invention relates to a liquid cosmetic composition and a powdery cosmetic composition having improved product characteristics such as useability and stability, and effectively preventing skin roughening, due to the formulation of an inclusion product with HACD.

2. Description of the Related Art

Cosmetic compositions well known in the art such as cosmetic water, lotions, beauty lotion, etc. contain, in addition to a main component such as water or alcohol, various slightly water-soluble components such as oils and fats, physiologically active substances, UV-ray absorbers, antiphlogistics, and perfumes, formulated by utilizing a small amount of a surfactant for enhancing the useability and utility thereof.

Also, powdery cosmetic compositions used for an amelioration of skin sores due to a sunburn contain a slightly water-soluble component such as a skin activator and a drug in a water-soluble powdery base.

In cosmetic compositions, it is well known that a quantitative limitation will arise when formulating the above-mentioned slightly water-soluble components, and therefore, for example, when solubilizing slightly water-soluble components in large

- 2 -

amounts and in a uniform state, a large amount of ethanol or a polyol must be formulated, but the formulation of these components cause the problem of skin irritation. Also, if the slightly water-soluble components are not uniformly dissolved, a problem arises in that the cosmetic composition becomes turbid or loses its transparency, to thereby reduce its commercial value. Further, the slightly water-soluble components formulated also react with the other components, to thereby cause a deterioration and decomposition of the other components.

To solve these problems, Japanese Unexamined Patent Publication (Kokai) No. 61-227517 discloses a technique by which the above-mentioned components are formulated by utilizing the inclusion action of a cyclodextrin polymer. But, according to this method, since a cyclodextrin polymer per se having a constant quality is difficult to obtain, and because the polymer is viscous and slightly soluble in water, satisfactory results could not be obtained.

Also, in powdery cosmetic compositions of the type which are commercially available in powdery form, to which water is added before use, since this type of cosmetic is solubilized between the palms while using only of a small amount of water, the slightly water-soluble components are not uniformly dissolved, and therefore, a coarse feeling remains during use and the pharmacological effect thereof is not fully exhibited. As means for solving these problems, attempts have been made to increase the amount of the surfactant or to utilize the inclusion action of a cyclodextrin polymer, but good results could not be obtained for the same reason as mentioned above. Also, none of the cyclodextrin polymers provides a satisfactory skin roughening prevention effect.

Recently, there is a trend toward using a base makeup on the skin, primarily directed to beautifying

- 3 -

the skin, and thus there is a need for a cosmetic composition which effectively prevents skin roughening and retains the preferable characteristics of a cosmetic composition.

5 SUMMARY OF THE INVENTION

Accordingly, the objects of the present invention are to eliminate the above-mentioned disadvantages of the prior art and to provide a cosmetic composition having a high degree of safety, an excellent solubility, useability, and stability, and providing an effective
10 skin roughening prevention, by utilizing the inclusion action of HACD, which can be readily solubilized without using a large amount of a solubilizing agent such as ethanol or a polyol.

15 Other objects and advantages of the present invention will be apparent from the following description.

In accordance with the present invention, there is provided a cosmetic composition comprising an inclusion
20 product having a slightly water-soluble component with a hydroxyalkylated cyclodextrin formulated therein.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

The HACD used in the present invention is a cyclodextrin well known in the art as a cyclic
25 oligosaccharide, having a hydroxyalkyl group introduced into the hydroxyl group thereof.

As the hydroxyalkyl group, substituents such as hydroxymethyl, hydroxyethyl, hydroxypropyl, and hydroxybutyl can be used, and as a result of the
30 reactions of these substituents, an HACD such as hydroxymethyl cyclodextrin, hydroxyethyl cyclodextrin, hydroxypropyl cyclodextrin, and hydroxybutyl cyclodextrin can be obtained.

Cyclodextrin (hereinafter abbreviated as "CD") is
35 known to include CD having α , β and γ structures, (hereinafter abbreviated as " α -CD", " β -CD" and " γ -CD"), depending on the difference in the glucose number, and

- 4 -

in the present invention, one or more of these cyclodextrins can be used after hydroxyalkylation. Also, a starch decomposed product containing α , β , and γ CD's can be used at the same time.

5 These HACD's have an excellent solubility in hydrophilic solvents such as water, compared with the CD of the prior art as shown in Table 1, and in the present invention any desired amount of a slightly water-soluble component can be formulated in a cosmetic composition by
10 utilizing this property and the inclusion action of HACD.

Table 1
Solubilities of CD and HACD in Hydrophilic Solvents

Sample	Water	Water/Methanol		Water/Ethanol		Water/Acetone	
		50/50	0/100	50/50	0/100	50/50	0/100
β -CD	1.85	0.3	-	1.3	-	2.6	-
Hydroxyethyl β -CD	57	280	75	250	<1	230	<1
Hydroxypropyl β -CD	115	87	93	162	<1	75	<1
Hydroxypropyl CD mixture	337	325	<1	300	<1	285	<1

Remarks: Numerical values in Table indicate grams dissolved per 100 ml of solvent

- 6 -

Of these HACD's, when the cost, production ease, useability and water solubility are taken into consideration, preferably, hydroxyethylated CD or hydroxypropylated CD, more preferably hydroxyethylated β -CD or hydroxypropylated β -CD is used, but the present invention is by no means limited thereto.

Also, the hydroxyethylated CD or hydroxypropylated CD are produced as mixtures of α , β and γ CD's, but such a mixture or an isolated product of an α , β and γ hydroxypropylated CD can be used.

Several processes for preparing HACD are known in the prior art, and an example thereof is shown below.

A 100 g amount of β -CD (manufactured by Nihon Shokuhin Kako, tradename: Seldex N) was dissolved in 150 ml of a 20% aqueous NaOH, 50 ml of propylene oxide was gradually added dropwise while maintaining the temperature at 30°C, and the reaction was continued for 20 hours under stirring. After completion of the reaction, the pH was adjusted to 6.0 with hydrochloric acid, and the product placed in a dialyzing film tube to effect desalting under running water for 24 hours. Then the product was dried by a lyophilizer, to give about 90 g of a propylated β -cyclodextrin. The degree of substitution of the hydroxypropylated β -cyclodextrin was found to be 5.1.

The slightly water-soluble component to be used in the present invention means a cosmetic component which is substantially insoluble or only slightly soluble in water. Specific examples include, for example, natural animal and vegetable oils and fats such as macadamia nut, evening primrose oil, olive oil, mink oil, jojoba oil, lanolin, and squalene; hydrocarbons such as fluid paraffin, squalane; waxes such as paraffin wax, whale wax, beeswax, candelilla wax, and carunauba wax; higher alcohols such as cetanol, isocetanol, stearyl alcohol, and isostearyl alcohol; higher fatty acids such as myristic acid, palmitic acid, stearic acid, behenic

- 7 -

acid, isostearic acid, oleic acid, linolenic acid, and linoleic acid oxyacid; fatty acid esters such as isopropyl myristate, isopropyl palmitate, isopropyl isostearate, and glyceryl 2-ethylhexanoate; polar oils
5 such as diethylene glycol monopropyl ether, polyoxyethylene polyoxypropylene pentaerythritol, polyoxypropylene butyl ether, and ethyl linoleate; and silicone oils; and the like.

Further, there may be included vitamins and
10 vitamin-like acting substances such as vitamin A, vitamin D, vitamin E, tocopherol acetate, and ascorbate, and g-oryzanol and folic acid; hormones such as estradiol benzoate, estradiol valerate, ethynyl estradiol, prostaglandin, and testosterone propionate;
15 UV-ray absorbers such as benzophenone, 4-t-butyl-4'-methoxydibenzoylmethane, glyceryl dimethoxycinnamate ethylhexanoate, p-aminobenzoate, octyl p-methoxycinnamate, and phenyl salicylate; antiphlogistics such as L-menthol and camphor; preservatives such as ethyl
20 p-benzoate, propyl p-benzoate, and butyl p-benzoate; and sterilizers such as glycyrrhizinic acid, triclosane, and dibutylhydroxy toluene. Also, there can be included oil-soluble dyes such as Oil Red, Naphthol Yellow, Tartrazine, and Puplika; single substance perfumes such
25 as linalool, linalyl acetate, limonene, citral, methylionone, benzyl acetate, methyldehydrojasmonate, phenylethyl alcohol, musk ketone, santalol, a-hexylcinnamicaldehyde, TEC, and citronelol, and composed perfumes comprising mixtures thereof; essential
30 oils derived from natural animal and vegetable oils and fats, and galenicals.

The property of these substances may be either liquid or crystalline, and they may be used either alone or in the form of a mixture.

35 When preparing the inclusion product using such a slightly water-soluble component and HACD, the well-known methods as described below may be employed.

- 8 -

To an aqueous 20 to 60% by weight of HACD is added and mixed 0.01 to 0.15 part by weight of the required component as mentioned above per 1 part by weight of HACD, followed by stirring and mixing at 20 to 50°C.
5 The stirring may be carried out at 50 to 3000 rpm, and the inclusion reaction time is 2 to 8 hours.

The inclusion product thus obtained is solubilized or emulsified in the aqueous solution, and can be used as such, but the aqueous solution can be also finely
10 powdered by a treatment such as freeze drying or spray drying before use. In the liquid or non-powdery cosmetic composition according to the first embodiment of the present invention, a formulation is possible in either the the reaction mixture or the finely powdered
15 state.

The content of the above-mentioned inclusion product in the liquid cosmetic composition of the present invention, which is influenced by the kind of slightly water-soluble component included and the type
20 of cosmetic composition, is preferably up to 25% by weight, more preferably 0.5 to 15% by weight of the entire cosmetic composition in the finely powdered state. Although it is technically possible to formulate an amount of more than 25% by weight, a drawback arises
25 in that the cosmetic composition becomes sticky during use.

The liquid or non-powdery cosmetic composition of the first embodiment of the present invention, in addition to the above components, can further formulate
30 other conventional cosmetic components corresponding to the commercial product feature, for example, humectants such as hyaluronic acid; water-soluble drugs such as vitamin C; viscosity controllers; pH controllers; preservatives; sterilizers, antioxidants; perfumes; and
35 dyes.

In the powdery cosmetic composition of the second embodiment of the present invention, the same inclusion

product as used in the cosmetic composition of the first embodiment of the present invention can be used, but since the dried product form is maintained until immediately before use, preferably the inclusion product of HACD is added in the dry state to other water-soluble base components. The amount of the inclusion product of HACD in the powdery cosmetic composition is preferably up to 50% by weight, more preferably 1.0 to 40%, of the entire powdery cosmetic in the finely powdered state. Although no substantial problem arises with an amount of more than 50% by weight, the cosmetic does become sticky during use.

In the powdery cosmetic composition of the second embodiment of the present invention, soluble powder bases such as D-mannitol and lactose other than the inclusion product of HACD, and corresponding to the commercial product feature, other cosmetic components, for example, humectants such as hyaluronic acid; water-soluble drugs such as vitamin C; viscosity controllers; pH controllers; preservatives; sterilizers; antioxidants; perfumes; and dyes can be also formulated.

According to the present invention, slightly water-soluble components which can be limitedly formulated in the prior art can be formulated in any desired amount without the aid of alcohol, whereby liquid or powdery cosmetic compositions are obtained which have an excellent transparency, stability, safety, useability, and effectively prevent skin roughening. Also, in the present invention, since the slightly water-soluble substance is included with HACD, other components in the liquid or powdery cosmetic compositions are not affected thereby, and thus a long term stabilization is obtained.

Further, in the second aspect of the present invention, an effect is obtained whereby the powdery cosmetic composition is completely dissolved with a small amount of water during use.

- 10 -

EXAMPLES

The present invention will now be further illustrated in detail by, but is by no means limited to, the following Examples, wherein "parts" in the formula-
 5 tion are all by weight unless otherwise noted.

Example 1: Humectant Cosmetic WaterPreparation 1

	(1) Hydroxypropylated CD mixture	2.5	parts
	(2) Macadamia nut oil	0.5	"
10	(3) Vitamin E acetate	0.05	"
	(4) Deionized water	40	"

Preparation 2

	(1) Deionized water	39.729	parts
	(2) Sorbitol	5	"
15	(3) 1,3-Butylene glycol	12	"
	(4) Lactic acid	0.02	"
	(5) Sodium lactate	0.1	"
	(6) Monoammonium glycyrrhizinate	0.1	"
	(7) Dye	0.001	"

Preparation method

The hydroxypropylated CD mixture of (1) (2.5 g) was dissolved in 40 g of deionized water, (2) and (3) were added thereto, and the mixture was stirred to obtain the Preparation 1 containing the inclusion product of the
 25 hydroxypropylated CD mixture.

Then the above Preparation 1 was added to the above Preparation 2 to obtain a humectant cosmetic water having a drug stably formulated therein.

When the hydroxypropylated CD mixture was omitted
 30 from the above-described composition, the resultant cosmetic water did not have a good stability, in that the oil components were separated from the drug over a lapse of time.

Example 2: Systemic LotionPreparation 1

	(1) Hydroxypropylated β -CD	5.0	parts
	(2) Benzophenone	0.05	"

- 11 -

	(3)	4-t-Butyl-4'-methoxy-dibenzoylmethane	0.01	"
	(4)	Glyceryl dimethoxycinnamate ethylhexanoate	0.01	"
5	(5)	Deionized water	2.0	"

Preparation 2

	(1)	Deionized water	49.8699	parts
	(2)	PEG 400	1.0	"
	(3)	Hinokitiol	0.01	"
10	(4)	Luffa extract	1.0	"
	(5)	Iris extract	1.0	"
	(6)	Modified 95% ethanol	40.0	"
	(7)	Perfume	0.05	"
	(8)	Dye	0.0001	"

15 Preparation method

To a solution of (1) dissolved in deionized water were added UV-ray absorbers (2), (3) and (4), followed by stirring, to prepare the Preparation 1 containing the inclusion product of hydroxypropylated β -CD.

20 Then the above Preparation 1 was added to the Preparation 2 to obtain a systemic lotion having UV-ray absorbers stably formulated therein.

When the hydroxypropylated β -CD was omitted from the above-described composition, lotion had a poor
25 stability, in that the UV-absorbers were separated over a lapse of time.

Example 3: EssencePreparation 1

	(1)	Hydroxypropylated α -CD	10	parts
30	(2)	Evening primrose oil	0.2	"
	(3)	α -Tocopherol	0.05	"
	(4)	Deionized water	20	"

Preparation 2

	(1)	Deionized water	59.26	parts
35	(2)	Dipropylene glycol	5	"
	(3)	Maltitol	5	"
	(4)	Aspartic acid	0.04	"

- 12 -

(5)	L-Arginine	0.1	"
(6)	Sodium hexametaphosphate	0.05	"
(7)	Carboxyvinyl polymer	0.2	"
(8)	Perfume	0.1	"

5 Preparation method

To a solution of (1) dissolved in deionized water were added (2) and (3), followed by stirring, to prepare the Preparation 1 containing the inclusion product of the hydroxypropylated α -CD.

10 Then the above Preparation 1 was added to the Preparation 2 to obtain an essence having an antioxidant stably formulated therein.

15 When the hydroxypropylated α -CD was omitted from the above-described composition, it was nonuniform and had a poor stability, in that the oil component was separated from the antioxidant over a lapse of time.

Example 4: Cosmetic Water

Preparation 1

(1)	Hydroxypropylated γ -CD	1.0	part
(2)	Benzophenone	0.05	"
(3)	Composed perfumes (limonene, phenylethyl alcohol, citronerol, etc.)	0.1	"
(4)	Deionized water	3	"

25 Preparation 2

(1)	Deionized water	82.0948	parts
(2)	Glycerol	1.0	"
(3)	1,3-Butylene glycol	2.0	"
(4)	Lactic acid	0.005	"
(5)	Sodium lactate	0.2	"
(6)	Monoammonium glycyrrhizinate	0.05	"
(7)	Aloe extract	0.5	"
(8)	Modified 95% ethanol	10.0	"
(9)	Dye	0.0002	"

35 Preparation method

To a solution of (1) dissolved in deionized water were added (2) and (3), to prepare the Preparation 1

- 13 -

containing the inclusion product of the hydroxy-propylated γ -CD.

Then the above Preparation 1 was added to the Preparation 2 to obtain a cosmetic water having a
5 UV-absorber and a perfume stably formulated therein.

Example 5: Astringent

Preparation 1

	(1) Hydroxyethylated β -CD	5.0 parts
10	(2) Composed perfume (Linanol, Methyldehydrojasmonate, Muskketone, Valinine, etc.)	0.03 "
	(3) 4-t-Butyl-4'-methoxy-dibenzoylmethane	0.01 "
	(4) Naphthol yellow	0.01 "
15	(5) Deionized water	20 "

Preparation 2

	(1) Deionized water	57.57 parts
	(2) Dipropylene glycol	2.0 "
	(3) Citric acid	0.03 "
20	(4) Sodium citrate	0.05 "
	(5) Zinc sulfophenolate	0.2 "
	(6) Modified 95% ethanol	15.0 "
	(7) Methyl p-benzoate	0.1 "

Preparation method

25 To a solution of (1) dissolved in deionized water were added (2), (3), and (4), followed by stirring, to obtain the Preparation 1 containing the inclusion product of the hydroxyethylated β -CD.

Then the above Preparation 1 was added to the
30 Preparation 2 to obtain an astringent having a UV-ray absorber, perfumes, and a dye stably formulated therein.

Example 6: Nonalcoholic Cosmetic Water

Preparation 1

	(1) Hydroxymethylated β -CD	10 parts
35	(2) Mentol	3 "
	(3) Perfume (Rose oil)	0.01 "
	(4) Deionized water	20 "

- 14 -

Preparation 2

	(1)	Deionized water	56.46	parts
	(2)	Dipropylene glycol	10	"
	(3)	Citric acid	0.03	"
5	(4)	Sodium citrate	0.05	"
	(5)	Methyl p-benzoate	0.1	"
	(6)	Phenoxyethanol	0.3	"
	(7)	Dye	0.05	"

Preparation method

10 To a solution of (1) dissolved in deionized water were added (2) and (3), followed by stirring, to prepare the Preparation 1 containing the inclusion product of the hydroxymethylated β -CD.

15 Then the above Preparation 1 was added to the Preparation 2 to obtain a nonalcoholic cosmetic water containing an antiphlogistic and a perfume stably formulated therein.

Example 7: Powder Containing Cosmetic WaterPreparation 1

20	(1)	Hydroxybutylated β -CD	6.0	parts
	(2)	l-Mentol	0.3	"
	(3)	Camphor	0.5	"
	(4)	Perfume	0.15	"
	(5)	Deionized water	15	"

Preparation 2

25	(1)	Deionized water	69.65	parts
	(2)	Glycerol	1.0	"
	(3)	Asparagine	0.05	"
	(4)	Modified 95% ethanol	5.0	"
30	(5)	Zinc	1.5	"
	(6)	Kaolin	0.5	"
	(7)	Methyl p-benzoate	0.05	"
	(8)	Clay mineral	0.3	"

Preparation method

35 To a solution of (1) dissolved in deionized water were added (2), (3), and (4), followed by stirring, to prepare the Preparation 1 containing the inclusion

- 15 -

product of the hydroxybutylated β -CD.

Then the above Preparation 1 was added to the Preparation 2 to obtain a powder containing a cosmetic water having an antiphlogistic and a perfume formulated stably therein.

Example 8: White Powder

Preparation 1

	(1)	Hydroxypropylated CD mixture	10	parts
	(2)	Benzophenone	0.1	"
10	(3)	4-t-Butyl-4'-methoxy-dibenzoylmethane	0.05	"
	(4)	Glyceryl dimethoxycinnamate ethylhexanoate	0.05	"
	(5)	Deionized water	10	"

Preparation 2

	(1)	D-Mannitol	64.77	parts
	(2)	L-Ascorbic acid	3.0	"
	(3)	Dipalmityl ascorbate	12.0	"
	(4)	Monoammonium glycerrhizinate	0.05	"
20	(5)	Riboflavin	0.02	"

Preparation method

To a solution of (1) dissolved in deionized water were added (2), (3), and (4), followed by stirring, to prepare the Preparation 1 containing the inclusion product of the hydroxypropylated CD mixture. The Preparation 1 was lyophilized to form a powder thereof.

Then the powder of the above Preparation 1 was added to the Preparation 2 to obtain a white powder having a UV-absorber stably formulated therein.

Example 9: Essence Powder

Preparation 1

	(1)	Hydroxypropylated CD mixture	8	parts
	(2)	Evening primrose oil	0.1	"
	(3)	α -Tocopherol	0.05	"
35	(4)	Hyaluronic acid	0.001	"
	(5)	Deionized water	12	"

Preparation 2

- 16 -

(1) Lactose 79.849 parts

Preparation method

To a solution of (1) and (4) dissolved in deionized water were added (2) and (3), followed by stirring, to prepare the Preparation 1 containing the inclusion product of the hydroxypropylated CD mixture. The Preparation 1 was spray dried to form a powder thereof.

Then the powder of the above Preparation 1 was added to the Preparation 2 to obtain an essence powder having an oil component, an antioxidant, and a humectant stably formulated therein.

Example 10: Suncare Powder

Preparation 1

	(1) Hydroxypropylated α -CD	4.0 parts
15	(2) Benzophenone	0.05 "
	(3) Deionized water	6 "

Preparation 2

	(1) Hydroxypropylated β -CD	4.0 parts
20	(2) 4-t-Butyl-4'-methoxy-dibenzoylmethane	0.02 "
	(3) Glycerol dimethoxycinnamate ethylhexanoate	0.05 "
	(4) p-Aminobenzoic acid ester	0.5 "
	(5) Deionized water	7 "

Preparation 3

25	(1) D-mannitol	77.349 parts
	(2) 2-Hydroxy-4-methoxybenzo-phenone-5-sodium sulfonate	1.0 "
	(3) Uracanic acid	0.001 "
30	(4) Monoammonium glycyrrhizinate	0.03 "

Preparation method

The hydroxypropylated α -CD was dissolved in deionized water and (2) was added to the resultant solution, followed by stirring, to form a UV-ray absorber composite solution, which was then dried to prepare the Preparation 1.

Then the hydroxypropylated β -CD was dissolved in

- 17 -

deionized water, and (2), (3) and (4) of the Preparation 2 were added to the resultant solution to form a UV-ray absorber composition solution, which was dried to prepare the Preparation 2.

5 Further, (1), (2), (3) and (4) of the Preparation 3 were mixed to prepare the Preparation 3, and finally, the Preparation 1 and the Preparation 2 were added to the Preparation 3 to obtain a suncare powder having UV-ray absorbers stably formulated therein.

10 Example 11: Action against Skin Roughening

Sample of the inclusion products having composed perfume included in CD prepared by dissolving 5 g of various CD or CD derivatives in 95 g of deionized water and adding 0.01 g of a composed perfume (linalol, 15 linalol acetate, lavender oil, santalol, coumarine, etc.), followed by stirring, were used to determine the skin roughening prevention effect.

The hair at the back of 3 Hartley-strain guinea pigs weighing 700 to 850 g was cut with a pair of 20 clippers, followed by a of depilation treatment. On day 4 after the hair was removed, 25 ml of an aqueous 3% sodium alkylbenzenesulfonate (LAS) solution was continuously applied once per day on 4 sites of the skin to which the depilation treatment was applied to cause 25 skin roughening. Further, at the same sites, after coating the LAS solution, 50 ml of each of the above-mentioned samples was applied once per day.

The changes in the state of the skin on day 7 after the initiation of the LAS aqueous solution application 30 were evaluated with regard to erythema and exfoliation according to the standards shown below; these are average values. Also, the totals of the respective average values of erythema and exfoliation are shown as the skin roughening score. The results are shown in 35 Table 2.

Evaluation of Erythema

- 18 -

<u>Score</u>	<u>Change</u>
0	No change
0.5	Rubor slightly recognized
1	Rubor recognized from state of skin surface
2	Rubor and edema recognized from state of skin surface
3	Rubor, edema and erosion recognized from state of skin surface
4	Rubor, edema, erosion and ulcer recognized from state of skin surface

Evaluation of exfoliation

<u>Score</u>	<u>Change</u>
0	No change
0.5	Slight drying, but no exfoliation
1	Slight exfoliation
2	Moderate exfoliation
3	Heavy exfoliation

- 19 -

Table 2

Sample	Erythema	Exfoli- ation	Skin roughening score (Total)	Remark
Hydroxypropyl- β -CD	0.2	0.6	0.8	Invention
α -CD	0.7	2.0	2.7	Comparison
β -CD	0.9	0.7	1.6	"
γ -CD	0.5	1.1	1.6	"
β -CD-polymer	1.1	1.2	2.3	"

15 Although the skin roughening prevention effect is
greater as the numerical value of skin roughening score
is lower, as apparent from Table 2, it is obvious that
the hydroxypropyl- β -CD of the present invention acts
very effectively to prevent skin roughening, compared
20 with the samples formulated for comparative purposes.

Example 12: Action on Corneum Turnover

To determine the effect on turnover of corneum, the
respective samples were applied in the same manner as in
Example 1 except that duncil chloride was applied on the
25 same sites before the application of the 3% aqueous LAS
solution of Example 11. From the change of a relative
fluorescent value with a lapse of time, the days when
only 50% of the original fluorescent value appeared were
calculated. The results are shown in Table 3. The
30 turnover of corneum is accelerated as the days in which
50% of the fluorescence disappears are lessened,
suggesting progress against the skin roughening.

Table 3

Sample	50% fluorescence disappearance days	Remark
Hydroxypropyl- β -CD	4.7	Invention
α -CD	3.8	Comparison
β -CD	3.4	"
γ -CD	3.7	"
β -CD-polymer	2.7	"

- 15 As apparent from Table 3, the hydroxypropyl- β -CD of the present invention is very effectively inhibits the corneum turnover, compared with the samples formulated for comparative purposes.

- 21 -

We Claim:

1. A cosmetic composition comprising an inclusion product having a slightly water-soluble component with a hydroxyalkylated cyclodextrin formulated therein.
2. A cosmetic composition as claimed in claim 1,
5 wherein the content of the inclusion product in the cosmetic composition is up to 25% by weight under the finely powdered state.
3. A cosmetic composition as claimed in claim 2,
10 wherein the content of the inclusion product in the cosmetic composition is 0.5 to 15% by weight.
4. A cosmetic composition as claimed in claim 1,
wherein the slightly water-soluble component is at least one compound selected from the group consisting of
15 natural animal and vegetable oils and fats,
hydrocarbons, waxes, higher alcohols, higher fatty acids, fatty acid esters, polar oils, and silicone oils.
5. A liquid cosmetic composition of claim 1.
6. A powdery cosmetic composition of claim 1.

Barrigar & Moss
81 Metcalfe Street, 7th Floor
Ottawa, Ontario
K1P 6K7
Agents for the Applicant